

A SPURIOUS CORRELATION IN AN INTERPOPULATION COMPARISON OF ATLANTIC SALMON LIFE HISTORIES¹

RANSOM A. MYERS

Department of Fisheries and Oceans, P.O. Box 5667, St. John's, Newfoundland A1C 5X1, Canada

AND

JEFFREY A. HUTCHINGS

Department of Biology, Memorial University of Newfoundland, St. John's,
Newfoundland A1B 3X9, Canada

Abstract. We tested two hypotheses concerning geographical variation in Atlantic salmon (*Salmo salar*) life histories: (1) mean age at first reproduction is positively correlated with growth rate at sea and (2) within-population variation in age at first reproduction first increases and then decreases with latitude. Data on growth and age at first reproduction were compiled from 41 populations in eastern North America. Data reliability was checked by a redetermination of ages based on scale examination. The proportion of fish that were incorrectly aged was small ($\approx 0.7\%$); however, aging errors were primarily of one kind; salmon that had previously spawned were misclassified as virgin fish of an older age class.

Growth rate at sea was found not to be positively correlated with age at maturation. Schaffer and Elson's (1975) positive correlation between growth and age at first reproduction can be attributed to a subtle statistical artifact caused by aging errors. We also found that within-population variation of age at maturation was not related to latitude.

We conclude that tests of life history theories should not assume constancy in life history traits, such as mortality, among populations.

Key words: Atlantic salmon; evolution; geographical variation; life history; reproductive strategies; *Salmo salar*; spawning age; spurious correlation; statistical artifact.

INTRODUCTION

Life history theory predicts that increased growth rate should lead to delayed maturation, all other factors remaining constant (Schaffer 1974a). However, high growth rates are usually associated with early maturation for fish and invertebrates (Alm 1959, Kanep 1974, Myers and Runge 1983, Policansky 1983, Hutchings and Morris 1985). This implies either that the theory is wrong or that there is a general covariation of other life history factors with growth rate such that the predictions of the theory are usually reversed.

A notable exception to the general pattern for fish was reported for the Atlantic salmon, *Salmo salar*, the anadromous form of which spends 1-3 yr at sea before returning to its natal stream to spawn (Jones 1959). Schaffer and Elson (1975) found a strong positive correlation between time spent at sea prior to maturation and growth rate at sea in North American populations of Atlantic salmon. This result supported their application of optimal life history theory to Atlantic salmon.

We investigated this inconsistency by examining the data upon which Schaffer and Elson (1975) based their correlation and by analyzing a more extensive data base to test their prediction. We found no evidence of

a correlation between growth rate at sea and sea age at maturation in Atlantic salmon. Furthermore, we could not verify their conclusion that within-population variation in age at maturation increases and then decreases with increasing latitude. Schaffer and Elson's (1975) result can be attributed to a subtle statistical artifact that is of general interest.

METHODS

For simplicity, one-sea-year (1SY) salmon refers here to fish that had spent 13-16 mo (encompassing one winter) at sea before maturing and returning to their natal streams to spawn. Similarly, 2SY and 3SY salmon had spent two and three winters at sea, respectively. Sea age at maturation refers to time spent at sea prior to spawning.

The rate of growth in length during the 1st 3 yr at sea decreases with age in Atlantic salmon (Allen et al. 1972). Hence, the unweighted linear regression of mean length on virgin sea age employed by Schaffer and Elson (1975) was inappropriate for this concave relationship. Growth rate at sea is more appropriately estimated as the difference between mean fork lengths of successive year classes. Fork length is the distance from the tip of the snout to the mid-fork of the tail.

Published data were compiled for salmon populations from rivers in eastern North America (Table 1).

¹ Manuscript received 30 October 1984; revised 11 September 1986; accepted 25 September 1986.

Age determinations had been based on scale analyses. Records from counting fences were used in preference to sport angling statistics when both kinds of data existed. We did not use commercial catch records and included data in our analysis only if information on the variation of length at a given age was available.

We examined the age-specific frequency distributions of length for populations in which the original scales were available for age verification. Data reliability was assessed by checking for outliers. We re-examined scales for individuals having lengths over two standard deviations (SD) from the population mean. Aging errors were primarily (over 70%) caused by interpretation of spawning marks as annuli (for details see Myers and Hutchings 1987); salmon that had previously spawned were often misclassified as virgin fish of an older sea age class. For example, the rate at which 1SY salmon (including those that had previously spawned) were misclassified as 2SY salmon in the Exploits River, Newfoundland, was 0.7%. These aging errors resulted in a systematic underestimation of the age-specific mean lengths for multi-sea-year salmon.

We used the following procedure to assess data reliability for populations in which original scales were not available. If length at a given age is normally distributed, then the expected percentage of individuals lying 2 SD below the population mean is 2.28. If more than twice this percentage of any age class was more than 2 SD below the population mean, the population was excluded from further analyses. This criterion excluded the data from the West River, Nova Scotia. Fifty percent of the fish classified as 2SY salmon in the West River were <65 cm in length, well within the size range characteristic of repeat-spawning 1SY fish. The robustness of our results was tested by including this river in the analysis.

Individual age-specific length data were not available for the Narraguagus River, Maine, and the Koksoak River in Ungava Bay, Québec. Means from these rivers were included because they represent Atlantic salmon near the northern and southern extremities of its range. Exclusion of these data did not affect the results.

RESULTS AND DISCUSSION

Growth at sea and sea age at maturation

Forty-one rivers satisfied our criteria for acceptance (Table 1). Our analyses (Fig. 1) failed to support a relationship between sea age at maturation and growth rate at sea ($r = 0.12$; $P > .05$; $n = 41$). Similar results were obtained by estimating growth rate at sea as the difference between mean lengths of fish that matured after 3 and 1 yr at sea ($r = -0.37$; $P > .05$; $n = 21$), and 3 and 2 yr at sea ($r = -0.12$; $P > .05$; $n = 21$). Exclusion of rivers with small sample sizes ($n < 5$ fish per age class) yielded similar correlations. The correlation was not significant when the suspect data from the West River were included.




FIG. 1. Mean sea age at maturation (from sample sizes in Table 1) plotted against growth rate at sea as measured by the difference in mean lengths of fish that matured after 2 yr at sea and those that matured after 1 yr at sea; $r = 0.12$; $P(r) > .05$. Symbols refer to Newfoundland (□); Nova Scotia (Δ); New Brunswick (◇); Québec (+); and Maine (×).

We also tested Schaffer and Elson's hypothesis within geographic regions by regressing mean sea age at maturation against growth rate at sea for each province. The regression for New Brunswick salmon was positive and nominally significant at the .05 level. That one test should appear to be significant of the 14 performed in our analysis was likely due to chance alone (the probability of obtaining at least one significant result by chance at the .05 level from 14 independent tests is 0.51). Furthermore, it is questionable whether New Brunswick comprises a reasonable geographic region for such an analysis; two of the rivers considered empty into the Bay of Fundy, whereas two empty into the Gulf of St. Lawrence.

Statistical artifacts

In the original Exploits River data, a 0.7% error in aging resulted in >50% of the alleged 2SY salmon being misclassified. Under what circumstances can a 0.7% classification error rate be responsible for a correlation that appears to be highly significant (e.g., $P < .001$)? If measurement errors associated with the variations in question are not independent, then even a relatively small error rate can have serious statistical consequences. In this case, age misclassifications affect estimated mean age as well as growth rate. This effect will be small if the number of observations in each class, for example age class, are approximately equal, but can become important if there is great disparity in sample sizes.

A simplified example will demonstrate this principle. Consider populations that consist only of 1SY and 2SY salmon. The mean sea age at maturation will be $1 + q$, where q is the proportion of the population returning after 2 yr at sea. Let e be the proportion of 1SY salmon misclassified as 2SY salmon. If all 1SY salmon are of length l_1 , and 2SY salmon of length l_2 , then growth as estimated by the apparent difference between the mean length of the two age classes is

TABLE 1. Relationship between length (cm) and number of years spent at sea (SY) for virgin Atlantic salmon in North America.

River	Latitude °N	Fork length* (sample size)			Collection method†	Year	Reference
		1SY	2SY	3SY			
Newfoundland							
Exploits	49.0	52.0 (846)	75.6 (10)		F	1963	Myers and Hutchings (1987)
Hunt	55.0	57.8 (8)	76.6 (24)		A	1980	Myers and Hutchings (1987)
North Harbour	47.0	55.1 (38)	76.3 (11)		F	1970, 1971, 1973, 1974	Lear and Day (1977)
Pinware	52.0	53.2 (748)	74.3 (142)		A	1980, 1981	Myers and Hutchings (1987)
Sand Hill	54.0	53.9 (3196)	73.0 (389)	98.1 (2)	A	1967, 1969-1974	Myers and Hutchings (1987)
Eagle	54.0	54.4 (302)	74.7 (9)		A	1971, 1979	Myers and Hutchings (1987)
Forteau	51.5	54.2 (353)	72.9 (11)		A	1979-1981	Myers and Hutchings (1987)
Highlands	48.0	53.3 (16)	73.4 (4)		F	1982	Myers and Hutchings (1987)
Humber	49.0	55.0 (297)	73.5 (9)	86.5 (17)	F	1967, 1976, 1979-1981	Myers and Hutchings (1987)
Little Codroy	47.5	54.1 (197)	75.1 (163)	89.6 (35)	F	1957-1963	Myers and Hutchings (1987)
Lomond	49.5	51.4 (108)	70.7 (7)		A	1975, 1978-1981	Myers and Hutchings (1987)
New Brunswick							
Miramichi	47.0	52.1 (8604)	72.8 (4016)	82.4 (48)	F	1971-1983	‡
Saint John	46.5	56.3 (583)	78.0 (474)		F	1977-1979	Myers and Hutchings (1987)
Big Salmon	45.0	53.2 (1654)	70.8 (139)		F	1965-1973	Jessop (1986)
Restigouche	48.0	53.2 (1488)	76.4 (701)	92.6 (335)	F	1972-1980	Randall (1984)
Nova Scotia							
LaHave	44.0	53.8 (68)	72.5 (28)		F	1982	Myers and Hutchings (1987)
Liscomb	45.0	52.3 (770)	72.4 (82)	91.7 (5)	F	1979-1982	Myers and Hutchings (1987)
Margaree	46.0	53.5 (17)	71.1 (23)	81.7 (5)	F	1973	Myers and Hutchings (1987)
North	46.0	54.6 (2)	72.3 (35)		F	1972	Myers and Hutchings (1987)
Philip	44.5	56.6 (22)	74.4 (167)	86.7 (34)	A	1978	Myers and Hutchings (1987)
St. Mary's	44.0	54.0 (60)	74.9 (20)	91.1 (9)	A	1974	Myers and Hutchings (1987)
Québec							
Bouleau	50.0	50.6 (15)	74.6 (20)		G	1972, 1973	Randall and Power (1979)
Pigou	50.0	53.1 (5)	72.7 (7)		G	1972, 1973	Randall and Power (1979)
Cap-Chat	49.0	56.3 (16)	78.9 (73)	93.4 (36)	A	1974, 1975, 1977	Myers and Hutchings (1987)
Corneille	50.0	53.3 (203)	70.0 (369)		A	1964-1968	Myers and Hutchings (1987)
Grande	48.0	54.9 (12)	74.4 (76)	90.9 (9)	A	1983	Myers and Hutchings (1987)
Koksoak	57.0	61.7 (221)	78.6 (253)		G	1980	Robitaille et al. (1984)
Leaf	58.5	62.4 (6)	79.7 (44)		G	1968	Lee and Power (1976)
Madeleine	49.0	51.5 (172)	75.2 (334)	92.1 (53)	A	1972, 1975-1979	Myers and Hutchings (1987)
Matamek	50.0	52.9 (20)	72.9 (6)		G	1974	Myers and Hutchings (1987)
Matane	49.0	54.8 (186)	71.9 (149)	90.3 (56)	A	1969	Myers and Hutchings (1987)
Matapédia	48.0	56.8 (66)	80.0 (354)	94.6 (171)	A	1973	Myers and Hutchings (1987)
Nabisipi	50.0	54.5 (61)	77.4 (1252)	87.6 (14)	G	1960-1962, 1963, 1965-1967	Myers and Hutchings (1987)
Pt. Saguenay	48.0	56.7 (182)	79.1 (39)	92.3 (83)	A	1978, 1979	Myers and Hutchings (1987)
Ste. Anne	49.0	56.1 (19)	80.8 (107)	95.8 (39)	A	1973, 1974	Myers and Hutchings (1987)
Ste. Marguerite	48.0	52.8 (63)	75.2 (70)	90.1 (75)	A	1969, 1976	Myers and Hutchings (1987)
St. Jean	50.0	53.0 (35)	75.9 (73)	89.3 (3)	A	1977	Myers and Hutchings (1987)
St. Jean (Gaspésie)	48.5	54.4 (60)	73.3 (157)	90.7 (11)	F	1978, 1979, 1983	Myers and Hutchings (1987)
Trinité	49.0	54.2 (319)	74.7 (21)		A	1981	Myers and Hutchings (1987)
Watshishou	50.0	53.3 (189)	70.4 (254)		A	1981-1983	Riley et al. (1984)
Maine							
Narraguagus§	44.5	54.8 (13)	69.8 (964)	81.7 (19)	F	1962-1966	Meister and Cutting (1967)

* Fork length = distance from tip of snout to mid-fork of tail.

† F = Counting fence; A = recreational angling; G = research gill nets within river/estuary.

‡ R. G. Randall, Canadian Department of Fisheries and Oceans, Moncton, New Brunswick.

§ Lengths given in Schaffer and Elson (1975) are total length, not fork length.

$$\frac{ql_2 + e(1-q)l_1}{q + e(1-q)} - l_1,$$

or equivalently

$$(l_2 - l_1) \frac{q}{q + e - eq} \quad (1)$$

The growth in length is greatly underestimated if q is small. If e and q are both small, then Eq. 1 can be approximated by

$$(l_2 - l_1) \frac{q}{q + e} \quad (2)$$

ACKNOWLEDGMENTS

Data from New Brunswick were provided by M. Chadwick, B. Jessop, and B. Randall; from Nova Scotia by L. Marshall and R. Semple; and from Newfoundland by E. Ash, B. Dempson, J. Gibson, H. Lear, M. O'Connell, R. Porter, J. Pratt, and D. Reddin. (All of the above are employees of the Canadian Department of Fisheries and Oceans.) Data for Québec were provided by Y. Côté and G. Ouellet, Ministère du Loisir, de la Chasse et de la Pêche, Gouvernement du Québec. Scale reading was checked by D. Reddin and B. Short, Department of Fisheries and Oceans, St. John's. We thank R. Haedrich, J. Rice, A. Rosenberg, and D. Schneider for comments on the manuscript.

LITERATURE CITED

- Allen, K. R., R. L. Saunders, and P. F. Elson. 1972. Marine growth of Atlantic salmon, *Salmo salar*, in the northwest Atlantic. *Journal of the Fisheries Research Board of Canada* 29:1373-1380.
- Alm, G. 1959. Connection between maturity, size and age in fishes. Institute of Freshwater Research Drottningholm. Report 40:5-145.
- Bell, G. 1980. The costs of reproduction and their consequences. *American Naturalist* 116:45-76.
- Gross, M. R. 1984. Sunfish, salmon, and the evolution of alternative reproductive strategies and tactics in fishes. Pages 55-75 in G. W. Potts and R. J. Wootton, editors. *Fish reproduction: strategies and tactics*. Academic Press, London, England.
- . 1985. Disruptive selection for alternative life histories in salmon. *Nature* 313:47-48.
- Hutchings, J. A., and D. W. Morris. 1985. The influence of phylogeny, size and behaviour on patterns of covariation in salmonid life histories. *Oikos* 45:118-124.
- Jessop, B. M. 1986. Atlantic salmon (*Salmo salar*) of the Big Salmon River, New Brunswick. Canadian Technical Report of Fisheries and Aquatic Sciences Number 1415.
- Jones, J. W. 1959. The salmon. Collins, London, England.
- Kanep, S. V. 1974. General features of the growth and sexual maturation of the whitefish (*Coregonus lavaretus*) in the USSR. *Journal of Ichthyology* 14:911-924.
- Lear, W. H., and F. A. Day. 1977. Analysis of biological and environmental data collected at North Harbour River, Newfoundland, during 1959-75. Canadian Department of Fisheries and Marine Services Technical Report Number 697.
- Lee, R. L. G., and G. Power. 1976. Atlantic salmon (*Salmo salar*) of the Leaf River, Ungava Bay. *Journal of the Fisheries Research Board of Canada* 33:2616-2621.
- Meister, A. L., and R. E. Cutting. 1967. A preliminary report of the composition of the spawning runs of Atlantic salmon, *Salmo salar*, in Maine rivers for the period 1962-66. International Commission for the Northwest Atlantic Fisheries, Redbook Part III:53-57.
- Myers, R. A. 1984. Demographic consequences of precocious maturation of Atlantic salmon, *Salmo salar*. *Canadian Journal of Fisheries and Aquatic Sciences* 41:1349-1353.
- . 1986. Game theory and the evolution of Atlantic salmon (*Salmo salar*) age at maturation. Pages 53-61 in D. Meerburg, editor. *Salmonid age at maturity*. Canadian Special Publication of Fisheries and Aquatic Sciences Number 89.
- . 1987. Redd superimposition and the evolution of salmonid age at maturation. *Oikos*, in press.
- Myers, R. A., and R. W. Doyle. 1983. Predicting natural mortality rates and reproduction-mortality trade-offs from fish life history data. *Canadian Journal of Fisheries and Aquatic Sciences* 40:612-620.
- Myers, R. A., and J. A. Hutchings. 1987. Sea age at maturity for Atlantic salmon, *Salmo salar*, in North America. Canadian Technical Report of Fisheries and Aquatic Sciences Number 1566.
- Myers, R. A., and J. A. Runge. 1983. Predictions of seasonal natural mortality rates in a copepod population using life-history theory. *Marine Ecology Progress Series* 11:189-194.
- Pauly, D. 1980. On the interrelationships between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks. *Journal du Conseil, Conseil International pour l'Exploration de la Mer* 36:175-192.
- Policansky, D. 1983. Size, age and demography of metamorphosis and sexual maturation in fishes. *American Zoologist* 23:57-63.
- Randall, R. G. 1984. Number of salmon required for spawning in the Restigouche River, N.B. Canadian Atlantic Fisheries Scientific Advisory Committee Research Document 84/16.
- Randall, R. G., and G. Power. 1979. Atlantic salmon (*Salmo salar*) of the Pigou and Bouleau rivers, Quebec. *Environmental Biology of Fishes* 4:179-184.
- Riley, S. C., A. T. Bielak, and G. Power. 1984. The Atlantic salmon stock of the Grand Watshishou River (Québec)—a historical perspective. *Naturaliste canadien* 111:219-228.
- Robitaille, J. A., Y. Côté, G. Hayeur, and G. Shoener. 1984. Particularités de la reproduction du saumon atlantique (*Salmo salar*) dans une partie du réseau Koksoak, en Ungava. *Rapport technique canadien des sciences halieutiques et aquatiques* 1313.
- Roff, D. A. 1984. The evolution of life history parameters in teleosts. *Canadian Journal of Fisheries and Aquatic Sciences* 41:989-1000.
- Schaffer, W. M. 1974a. Selection for optimal life histories: the effects of age structure. *Ecology* 55:291-303.
- . 1974b. Optimal reproductive effort in fluctuating environments. *American Naturalist* 108:783-790.
- Schaffer, W. M., and P. F. Elson. 1975. The adaptive significance of variations in life history among local populations of Atlantic salmon in North America. *Ecology* 56:577-590.
- Stearns, S. C. 1977. The evolution of life history traits: a critique of the theory and a review of the data. *Annual Review of Ecology and Systematics* 8:145-171.
- Stearns, S. C., and R. E. Crandall. 1984. Plasticity for age and size at sexual maturity: a life-history response to unavoidable stress. Pages 13-33 in G. W. Potts and R. J. Wootton, editors. *Fish reproduction: strategies and tactics*. Academic Press, London, England.